

***Tachysphex peruanus*, a New Species Related to  
*Tachysphex galapagensis* Williams  
(Hymenoptera: Sphecidae)**

WOJCIECH J. PULAWSKI

California Academy of Sciences, Golden Gate Park, San Francisco, California 94118.

---

*Abstract.*—*Tachysphex peruanus*, previously unknown, is a sister species of *Tachysphex galapagensis* and occurs in the coastal area of Peru.

---

A species not included in my revision (Pulawski, 1974) of Neotropical *Tachysphex* was found in material recently received for study. Relevant information is presented below. I sincerely thank Mary Ann Tenorio for taking the SEM picture and for drawing the distributional map.

***Tachysphex peruanus*, NEW SPECIES**

*Name derivation.*—Peruanus is a neo-Latin adjective derived from Peru, where the specimens have been collected.

*Diagnosis.*—Like other members of the *terminatus* species group, *peruanus* has a swelling behind each hindocellus combined with a flat, nonprominent labrum and punctate mesopleuron. Like *clarconis* W. Fox and *galapagensis* Williams, but unlike the remaining species of the group, the metapleuron of *peruanus* is simple (without carina or prominence in its upper part), and the metapleural flange is not expanded. Unlike all other species of the *terminatus* group, the clypeal lobe of the male is markedly angulate in *peruanus* (instead of truncate or arcuate), and the inner mandibular angle is simple (rather than dentate). Unlike *clarconis*, the vertex of *peruanus* is narrow (width to length ratio 1.3–1.4 in the female and 1.6–1.7 in the male instead of 1.9–2.2 and 2.1–2.6) and in the male the foretarsal rake is absent. Unlike *galapagensis*, the propodeal side is ridged in *peruanus* (nonridged or partly microridged in the former).

*Description.*—Mesopleural punctures more conspicuous posteriorly than in *galapagensis*, but less than in *clarconis*. Metapleural flange narrow. Upper metapleuron with a few simple, longitudinal ridges in front of propodeal spiracle, without prominence in its posterior part, without oblique carina beneath fore end of flange. Propodeum finely sculptured between dorsum and side, side ridged. Basal tooth of hindcoxa low, obtuse.

Vestiture partly obscuring integument between antennal socket and orbit. Setae length: on vertex about 1.5 times midocellar diameter in female and about 1.7 in male; about 1 midocellar diameter on scutum anterolaterally.

Head, thorax and legs black, except the following: mandible reddish yellow at about two-thirds of its length, and tarsal apex brown. Color of gaster variable,




Figure 1. *Tachysphex peruanus* species nova—clypeus of male.

gastral terga I–IV fasciate apically (fascia of tergum IV less conspicuous than the remaining ones). Wings weakly infumate.

♀: Clypeus width 2.8 times its midlength; bevel poorly defined; lip arcuate or very obtusely angulate, without median projection. Dorsal length of flagellomere I 2.4–2.5 times its apical width. Frons dull, microsculptured between punctures. Vertex width 1.3–1.4 times its length. Length 7.3–9.8 mm. Gaster all red or with dark, irregular spots.

♂: Mandibular inner margin nondentate (Fig. 1). Clypeus width 2.8 times its width; lip angulate, with free margin concave on each side of apex (Fig. 1). Dorsal length of flagellomere I 1.3–1.5 times its apical width. Frons punctatorugose. Vertex width 1.6–1.7 times its length. Gastral sterna pubescent throughout. Fore-femur notched. Foretarsomeres without preapical rake spines. Length 6.0–7.3 mm. Gaster differently colored in each specimen studied: all black, black basally (segments I–III) with the remaining segments red, and predominantly red (only segment I black).

*Discussion.* — The simple, unspecialized metapleuron of *clarconis*, *galapagensis*, and *peruanus* is a symplesiotypic character and therefore does not demonstrate close relationship. However, it does indicate that the three species are rather distinct from the remaining members of the group (in which the metapleuron is variously modified). Two derived characters are shared by *galapagensis* and *peruanus*, a narrow vertex and a reduced (some *galapagensis*) or absent foretarsal rake in the male, thus demonstrating their common origin. This is of great interest, since *peruanus* lives in an area from which ancestors of *galapagensis* most prob-




Figure 2. Geographic distribution of *Tachysphex galapagensis* Williams and *peruanus*, n. sp.

ably colonized the Galapagos Archipelago. However, each of the two species has at least one specialized character absent in the other (nonridged or barely ridged propodeal side in *galapagensis*, peculiar male clypeus and nondentate male mandible in *peruanus*), obviously precluding the latter from being the ancestor of the former. Clearly, the two are sister species, and must have evolved in parallel after their common ancestor colonized the Galapagos.

*Geographic distribution* (Fig. 2).—Coastal Peru, including foothills.

*Material examined.*—Holotype: ♀, Peru, Lambayeneque Province, Lambayeneque sand dunes, 2 Sep 1979, M. E. Irwin (1 ♂, University of California, Davis).

Paratypes (all from Peru): Lambayeneque Province: Same data as holotype (1 ♂, California Academy of Sciences; 1 ♂, University of California, Davis). Lima Province: Chosica, 2800 feet, 11 June 1914, H. S. Parish (1 ♂, Cornell University); Matucana, 17 May 1920, Cornell Univ. Expedition (1 ♀, Cornell University).

#### LITERATURE CITED

- Pulawski, W. J. 1974. A revision of the Neotropical *Tachysphex* Kohl (Hym., Sphecidae). *Polskie Pismo Ent.*, 44:3–80.

#### PUBLICATIONS RECEIVED

Trichoptera of the Area Platense. By Oliver S. Flint. *Biología Acuática* [Instituto de Limnología ILPLA], No. 2, 70 pp., 71 figs. Published by Instituto de Limnología "Prof. Dr. Raúl A. Ringuelet." Indicated as "noviembre 1982" issue. Available from Analía C. Paggi, Instituto de Limnología "Prof. Raúl A. Ringuelet," Casilla de Correo 55, 1923 Baerisso, Argentina (Bank drafts orders should be written in to the name of Analía C. Paggi). ISSN 0326-1638. Price US \$8 (postage fees included). Review copy received by PCES at CAS 31 May 1983.

New Zealand Butterflies. Identification and Natural History. By George W. Gibbs. William Collins Publishers Ltd., P.O. Box No. 1, Auckland. 207 pp., 63 figs., 197 colored "plates." Available from International Specialized Book Services, Inc., 10230 S.W. Parkway, Portland, Oregon 97225. First published 1980. Price \$45 hard cover. Review copy received by PCES at CAS 28 June 1983.

The Ecology of Aquatic Insects. Edited by Vincent H. Resh and David M. Rosenberg. Praeger Publishers, 521 Fifth Avenue, New York, New York 10175. x + 625 pp., figs. ISBN 0-03-059684-X. Publication date: March 1984. Price \$35 hard cover. Review copy received by PCES at CAS 2 April 1984.

Insect Ecology. Second Edition. By E. G. Matthews and R. L. Kitching with colour photographs by Densy Clyne. University of Queensland Press. xv + 211 pp., 14 figs., 12 illustrations, and 21 colored "plates." ISSN 0156-6695. Publication date given as 16 October 1984. Available from Queensland University Press, 5 South Union Street, Lawrence, Massachusetts 01843. Price \$15.95 paperbound. Review copy sent to PCES at CAS 7 May 1985.

The Pleasures of Entomology. Portraits of Insects and the People Who Study Them. By Howard Ensign Evans. Illustrated by Peter Eades. Smithsonian Institution Press. 238 pp., figs. ISBN 0-87474-421 (alk. paper). Paperbound. Price not indicated. Review copy received by PCES at CAS 31 May 1985. Available from Smithsonian Institution Press, Publications Distribution Section, 1111 North Capitol Street, N.E., Washington, D.C. 20560.—P. H. Arnaud, Jr., California Academy of Sciences, Golden Gate Park, San Francisco, California 94118.